

Please consider an early dose of the new corona vaccine.



The third dose of the COVID-19 vaccine can be given 6 months after the second dose. As speed is more important than type of the vaccine, once you receive your vaccination ticket, you should be vaccinated as soon as possible.

Q1

Why is the third vaccination necessary?

The effectiveness of the vaccine gradually declines over time, and it has been reported that a third dose of the vaccine can increase the effectiveness of the vaccine preventing infection and severe disease.



Q2

Does the third inoculation work for the Omicron strain?

The efficacy of Pfizer or Takeda/Moderna vaccines against Omicron strains in preventing severe illness requiring hospitalization has been reported to decrease over time, but, to recover after the third dose.

※The British Government reports that the effectiveness in preventing Omicron strain infection was 44% after 25 weeks after the second vaccination, but recovered to 89% after 2 weeks after the third vaccination.



Q3

Is it safe to give the third vaccination with a different type of vaccine than the one given in the first and second vaccinations?

In Japan, Pfizer and Takeda/Moderna are approved for use regardless of the type of vaccine given in the first and second doses. Both vaccines have been shown to be sufficiently effective and safe.

Vaccine	Combination	Antibody
● → ● → ○ Pfizer - Pfizer - Moderna		
● → ● → ● Pfizer - Pfizer - Pfizer		

◎Those who have not received their first and second vaccinations should continue to receive them.

◎There is an Immunization Health Damage Relief System.

Ogaki City Call Center for the COVID-19 Vaccination
TEL: 0584-47-6101 (8:30~17:15, Open weekdays as well as weekends and holidays)



Read this QR code to go to the booking site.